

Waterbody 3a
data delivered by UDU

Year	2002	
	Nov	Dec
Month		
Sampling point	S	S
DO ₂ (mg l ⁻¹)	13.0	NP
Temp (°C)	8.0	NP
pH	7.6	6.8
Conductivity (µS)	NP	NP
Cyanobacterial genera	NO	<i>Pla</i>
Chlorophyll <i>a</i> (µg l ⁻¹)	NP	3.8
Phaeophytin (µg l ⁻¹)	NP	0
Microcystin sc [†]	NA	NA
Microcystin in [†]	NP	0
Microcystin ex [†]	NP	0
Anatoxin-a sc [‡]	NA	NA
Anatoxin-a in [‡]	NP	0
Anatoxin-a ex [‡]	NP	0
Cylindrospermopsin sc [‡]	NA	NA
Cylindrospermopsin in [‡]	NP	0
Cylindrospermopsin ex [‡]	NP	0

NOTES

Sampling point: A, abstraction point for drinking water, S, surface water

Cyanobacterial genera: NO, not observed; OG, Other cyanobacterial genera present; *Mic*, *Microcystis*; *Ana*, *Anabaena*; *Aph*, *Aphanizomenon*; *Cyl*, *Cylindrospermopsis*; *Pla*, *Planktothrix*.

Toxins: sc, scum; in, intracellular toxin, filtered water sample; ex, extracellular toxin, filtered water sample; NA, not available; NP, not performed.

Toxin scale (extracellular and intracellular): 0, below minimum detection limit (<0.20µg l⁻¹); 1, 0.21-0.99 µg l⁻¹; 2, 1.00-5.00 µg l⁻¹; 3, 5.01-20.00 µg l⁻¹; 4, 20.01-100µg l⁻¹; 5, >100µg l⁻¹. Toxin scale (scum): 0, below minimum detection limit (<0.10µg g⁻¹); 1, 0.11-0.99 µg g⁻¹; 2, 1.00-10.00 µg g⁻¹; 3, 10.01-100.00 µg g⁻¹; 4, >100µg g⁻¹.

Where multiple methods for toxin analysis of an individual sample have been used, the highest observed concentration is recorded.

[†], Microcystin-LR equivalents measured by high performance liquid chromatography (HPLC), protein phosphatase inhibition assay and/or microcystin ELISA.

[‡], Anatoxin-a and cylindrospermopsin measured by HPLC.